



Consiglio Nazionale delle Ricerche
Istituto di Scienze dell'Atmosfera e del Clima



ITALIAN CLIMATE OBSERVATORY “O. VITTORI” Mt. CIMONE GAW-WMO Global Station



SPRING 2015 REPORT



CNR - ISAC
National Research Council
Institute of Atmospheric Sciences and Climate
ITALY

AUTHORS: Rocco Duchi, Paolo Cristofanelli, Angela Marinoni, Francescopiero Calzolari, Fabrizio Roccato, Davide Putero, Tony Christian Landi, Maurizio Busetto, Ubaldo Bonafè, Michela Maione, Igor Arduini and Paolo Bonasoni.



Institute of Atmospheric Science and Climate – National Research Council of Italy
Via Gobetti 101
40129 I-Bologna

Spring report of the Italian Climate Observatory "O. Vittori" at Mt.Cimone – n.8

Issued: November 2015, rev. 1

The report is published quarterly. Download at: <http://www.isac.cnr.it/cimone/reports>

ISSN: 2283-9631

The research activities and the measurement programmes carried out at the Italian Climate Observatory

"O. Vittori" at Mt. Cimone are supported by:

*The research activities and the measurement programmes carried out at the Italian Climate Observatory
"O. Vittori" at Mt. Cimone are an Italian contribute to:*



MIUR

Ministry of Education, University and Research
Ministero dell'Istruzione, dell'Università e Ricerca



CNR

National Research Council of Italy
Consiglio Nazionale delle Ricerche



DTA

Earth and Environment Department

Dipartimento di Scienze del Sistema Terra e Tecnologie per l'Ambiente



ACTRIS

Aerosol, clouds and trace gases research infrastructure network
Rete di Infrastrutture per la ricerca su aerosol, nubi e gas in tracce



NEXTDATA

A national system for the retrieval, storage, access and diffusion of environmental and climate data from mountain and marine areas.

Un sistema nazionale per la raccolta, conservazione, accessibilità e diffusione dei dati ambientali e climatici in aree montane e marine.



SHARE

Stations at High Altitude for Research on the Environment
Stazioni di alta quota per la ricerca sull'ambiente



WDCGG

World Data Center for Greenhouse Gases

Centro Dati Mondiale per i Gas Serra

<http://ds.data.jma.go.jp/gmd/wdcgg/>



WDCA

World Data Center for Aerosol

Centro Dati Mondiale per gli Aerosol

<http://www.gaw-wdca.org/>



MACC

Monitoring Atmospheric Composition & Climate

<http://gmes-atmosphere.eu/>



SDS-WAS

WMO Sand and Dust Storm Warning Advisory and Assessment System

<http://sds-was.aemet.es/>



AGAGE

Advanced Global Atmospheric Gases Experiment

<http://agage.eas.gatech.edu/>

ACKNOWLEDGEMENTS: ISAC-CNR gratefully acknowledges the Italian Air Force – CAMM Monte Cimone for the fundamental logistic support at Monte Cimone.

RINGRAZIAMENTI: l'ISAC-CNR ringrazia l'Aeronautica Militare Italiana – CAMM Monte Cimone per il fondamentale supporto logistico fornito.

DISCLAIMER: the data, analyses and information provided in this document should be considered as provisional. Significant changes can occur after the final revision of data. ISAC-CNR and the funding Projects reject any responsibility for the improper use of the information provided here. Reproduction or diffusion of the document contents is not allowed without the agreement of ISAC-CNR.

DICHIARAZIONE DI NON RESPONSABILITA': i dati, le analisi e le informazioni contenute in questo rapporto sono da considerarsi preliminari. Variazioni, anche significative, potranno verificarsi a seguito della validazione finale dei dati. ISAC-CNR ed i progetti finanziatori declinano ogni responsabilità per l'uso improprio delle informazioni contenute in questo rapporto. La riproduzione o la diffusione di questo documento non sono permesse senza l'autorizzazione esplicita dell'ISAC-CNR.

List of contents

Foreword	6
Premessa	7
Monte Cimone GAW/WMO Global Station.....	8
La Stazione Globale GAW/WMO di Monte Cimone	9
List of parameters.....	12
Lista dei parametri.....	13
Summary.....	14
Sommario	15
Special events	20
Eventi speciali.....	21
Surface ozone	28
Carbon monoxide (NDIR).....	30
Nitrogen oxides	32
Sulphur dioxide.....	34
Black carbon	35
Aerosol light scattering coefficient.....	37
Aerosol number concentration (fine).....	39
Aerosol number concentration (coarse)	41
Air Temperature	43
Relative humidity.....	45
Atmospheric pressure	47
Wind speed and direction	49
Solar radiation (short-wave and UV-B).....	51
Bibliography.....	53
Research Projects	58
The ICO-OV staff	61

Foreword

This report summarizes the results concerning the atmospheric observations carried out during Spring 2015 at the Italian Climate Observatory “O. Vittori” (ICO-OV), managed by the Institute of Atmospheric Sciences and Climate (ISAC) of the National Research Council of Italy (CNR). This research infrastructure is part of the WMO/GAW global station of Monte Cimone together with the Meteorological Observatory of the Italian Air Force(GAW ID: CMN).

Firstly, we provide a brief description of the measurement site and running experimental programmes. Then, an overview of the atmospheric and weather conditions during Spring 2015 is provided, by considering:

- **surface ozone**
- **carbon monoxide**
- **nitrogen oxides**
- **sulphur dioxide**
- **black carbon**
- **aerosol scattering coefficient**
- **fine and coarse particles**
- **meteorological data (temperature, relative humidity, pressure, wind speed and direction)**
- **solar radiation and UV-B**

For each atmospheric parameter, we provide basic statistical information (minimum, maximum and average values) together with a comparison with the climatological reference for Mt. Cimone.

Then, we present a list of special events, together with a description of the adopted selection methodologies:

- **pollution transport**
- **mineral dust transport**
- **transport of air-masses from the stratosphere**

For each observed parameter, a specific paragraph presents:

- **the time series of the daily mean values (calculated basing on 30-minute aggregated values, if the daily data coverage of 75% has been achieved)**
- **a table reporting the basic statistical parameters (on a 30-minute basis)**
- **a comparison with the seasonal historical mean values: for each year, the Spring mean values are calculated by averaging data from 2015, March 1st to 2015, May 31th.**

Premessa

Questo rapporto riassume i risultati relativi alle osservazioni atmosferiche effettuate durante la primavera 2015 presso l'Osservatorio Climatico "O. Vittori" (ICO-OV) dell'Istituto di Scienze dell'Atmosfera e del Clima (ISAC) del Consiglio Nazionale delle Ricerche Italia (CNR). Questa stazione di ricerca è parte, insieme all'Osservatorio Meteorologico dell'Aeronautica Militare, della stazione globale WMO/GAW di Monte Cimone (GAW ID: CMN).

In questo Report viene innanzitutto fornita una breve descrizione del sito di misura e dei programmi di ricerca in atto.

Viene quindi fornita una panoramica delle condizioni atmosferiche e meteorologiche che hanno caratterizzato la stagione primaverile del 2015 considerando:

- **ozono superficiale**
- **monossido di carbonio**
- **ossidi di azoto**
- **anidride solforosa**
- **black carbon**
- **particolato fine e grossolano**
- **gas alogenati**
- **composti organici volatili**
- **dati meteorologici (temperatura, umidità relativa, pressione, velocità e direzione del vento)**
- **radiazione solare e UV- B**

Per ogni parametro atmosferico sono fornite informazioni statistiche di base (valori minimi, massimi e medi) ed un confronto con il riferimento climatologico dell'Osservatorio "O. Vittori" per Monte Cimone.

Successivamente viene presentata una lista di eventi "speciali" identificati con procedure opportunamente messe a punto e descritte.

- **trasporto di masse d'aria inquinate**
- **trasporto di polvere minerale**
- **trasporto di masse d'aria dalla stratosfera**

Per ogni parametro osservato uno specifico paragrafo presenta:

- **Le serie storiche dei valori medi giornalieri (calcolati basandosi su valori mediati di 30 minuti, se la copertura dei dati giornalieri del 75% è stata raggiunta)**
- **Una tabella con i parametri statistici di base (su un base di 30 minuti)**
- **Il confronto con i valori medi storici stagionali per ogni anno, considerando che i valori invernali sono calcolati come media dal 1 marzo 2015 al 31 maggio 2015.**

Monte Cimone GAW/WMO Global Station

The **Global Atmosphere Watch (GAW)** programme of WMO is a partnership involving the Members of WMO, contributing networks and collaborating organizations and bodies which provides reliable scientific data and information on the chemical composition of the atmosphere, its natural and anthropogenic change, and helps to improve the understanding of interactions between the atmosphere, the oceans and the biosphere.



A network of measurement stations is the backbone of the GAW programme. This network consists of GAW Global and Regional measurement stations with additional measurements from Contributing stations. Both Global and Regional stations are operated by their host countries, either by their National Meteorological Services or by other national scientific organizations. More than 80 countries actively host GAW stations.

Currently GAW coordinates activities and data from 30 Global stations, more than 400 Regional stations, and around 100 Contributing stations operated by Contributing networks



*Location of the 30 Global Stations
of the WMO/GAW programme*

Mt. Cimone is the only WMO/GAW Global Station in Italy

Global station name	Altitude (a.s.l.)	Country
Assekrem/Tamanrasset	2710 m	Algeria
Izaña	2372 m	Spain
Jungfraujoch	3580 m	Switzerland
Mauna Loa,	3397 m	United States
Monte Cimone*	2165 m	Italy
Mt. Kenya	3678 m	Kenya
Mt. Waliguan	3810 m	China
Nepal Climate Observatory – Pyramid*	5079 m	Nepal
Zugspitze/ Hohenpeissenberg	2962 m	Germany

*Managed by Italian Institutions

List of GAW/WMO high altitude global station (for more information: <http://gaw.empa.ch/gawsis/>)

La Stazione Globale GAW/WMO di Monte Cimone

Il programma **Global Atmosphere Watch (GAW)** dell'OMM coinvolge gli Stati Membri della OMM e diverse reti osservative, organizzazioni ed Istituzioni con lo scopo di fornire dati scientifici ed informazioni attendibili sulla composizione dell'atmosfera, sui cambiamenti naturali e dovuti alle attività umane, contribuendo a migliorare la conoscenza delle interazioni fra atmosfera, oceani e biosfera.



La spina dorsale del programma GAW è costituita dalla propria rete osservativa. Essa è formata da Stazioni Globali e Regionali, oltre che da Stazioni definite "Contributing" che forniscono informazioni addizionali. Le Stazioni Globali e Regionali sono gestite dai Paesi di appartenenza, sia attraverso i Servizi Meteorologici Nazionali che gli Enti Pubblici di Ricerca. Le Stazioni GAW sono ospitate in oltre 80 Paesi del mondo.

*Alla stato attuale il programma coordina
30 Stazioni Globali e oltre 400
Stazioni Regionali, oltre a 100 Stazioni
"Contributing".*

*Dislocazione delle 30 Stazioni Globali
del programma WMO/GAW*



Monte Cimone è l'unica Stazione Globale WMO/GAW in Italia

Nome	Quota (s.l.m.)	Paese
Assekrem/Tamanrasset	2710 m	Algeria
Izaña	2372 m	Spagna
Jungfraujoch	3580 m	Svizzera
Mauna Loa	3397 m	USA
Monte Cimone*	2165 m	Italia
Mt. Kenya	3678 m	Kenya
Mt. Waliguan	3810 m	Cina
Nepal Climate Observatory – Pyramid*	5079 m	Nepal
Zugspitze/ Hohenpeissenberg	2962 m	Germania

*Stazioni gestite da Istituzioni Italiane


Lista delle stazioni globali GAW/WMO in aree di alta quota (for more information: <http://gaw.empa.ch/gawsis/>)

Geographical location

Mt. Cimone ($44^{\circ}12' N$, $10^{\circ}42' E$, 2165 m a.s.l.) is the highest peak of the Northern Apennines, the border line of two different climatic regions: the continental Europe northwards and the Mediterranean Basin southwards.

The closest inhabited areas are small villages placed 15 km from and about 1100 m below the Observatory, whereas major towns (500000 inhabitants) are situated in the lowlands about 60 km away (Bologna, Firenze). The industrial areas are not closer than 40 km and 2 km lower. The closest roads with some traffic are 7 km far and 1 km lower. Forest of conifers and beech trees grow up to 1600 m, so that the Laboratory is above the timberline. Only some patches of vegetation are on the top of the mountain

Mt. Cimone is characterized by a completely free horizon for 360° and air masses originated in different areas can reach the station. In the following figure, the annual 48-hour catchment areas, (i.e. the areas from which the air masses come) is provided for ICO-OV (EU-Project GEOMON).



The catchment area of the site for the 48-hour integration time backwards in time is given by the intersection of the topography with the volume containing the largest residence time densities and comprises 50% of the total residence time.

Il "bacino di raccolta" delle masse d'aria a 48 ore è ottenuto considerando le aree geografiche sopra le quali si totalizza almeno il 50% del tempo di residenza totale delle masse d'aria durante il trasporto verso Monte Cimone.

For more information:
<http://geomon.empa.ch/>.

The atmospheric observations carried out at Monte Cimone can be considered representative of the free tropospheric conditions of the Mediterranean Basin/South Europe. Only during the warm periods of the year, the measurement site can be affected by transport of air masses from PBL (planetary boundary layer).

Le osservazioni di composizione dell'atmosfera condotte a Monte Cimone possono essere considerate rappresentative delle condizioni di fondo della libera troposfera del bacino del Mediterraneo e del Sud Europa. Solo durante i mesi caldi, i processi convettivi possono favorire il trasporto di masse d'aria dallo strato limite planetario (PBL).

Posizione geografica

Monte Cimone (44°12' N, 10°42' E, 2165 m s.l.m.) è la cima più alta dell'Appennino Settentrionale, la linea di confine tra due diverse regioni climatiche: l'Europa continentale a Nord ed il bacino del Mediterraneo a Sud.

Le zone abitate più vicine sono piccoli paesi a circa 15 km di distanza e 1100 m di più in basso rispetto all'Osservatorio, mentre le città più grandi (Bologna, Firenze) sono situate in pianura a circa 60 km di distanza. Non vi sono importanti aree industriali nel raggio di circa 40 km. Le strade trafficate più vicine distano circa 7 km (1 km di quota più in basso). Boschi di conifere e faggi crescono fino a 1600 m. Nei pressi della cima si trovano prati e zone rocciose.

Monte Cimone è caratterizzato da un orizzonte completamente libero e quindi le masse d'aria possono raggiungere il sito di misura senza incontrare ostacoli orografici. Nella pagine precedente viene mostrata la media annuale del "bacino di raccolta" delle masse d'aria che nel giro di 48 ore sono arrivate a Mt. Cimone (EU-Project GEOMON).

NEWS

New selection criteria for the preliminary detection of stratospheric intrusion and pollution transport events

To perform a smart but reliable preliminary identification of free tropospheric/stratospheric air-mass transport and atmospheric pollution events at CMN, new selection methodologies are used starting with the current report.

To identify "dry" free tropospheric/stratospheric air-masses at CMN, we used a selection criterion used at the Zugspitze GAW/WMO Global Station – Germany (see Trickl et al., 2010): (i) RH continuously lower than 60%, (ii) RH minima lower than 30%; (iii) air-masses with potential vorticity higher than 1.6 pvu.

With the aim of identifying the presence of polluted air-masses at CMN, we selected the measurement periods for which at least two "primary" pollutants (CO, NO, BC, NO₂) showed average daily values higher than the monthly climatological mean value.

Nuovi criteri di selezione adottati per l'identificazione preliminare degli eventi di intrusione stratosferica e di trasporto di masse d'aria inquinate

Per semplificare e rendere più robusta l'identificazione preliminare degli eventi di trasporto di masse d'aria provenienti dalla libera troposfera/bassa stratosfera e di masse d'aria inquinate, a partire da questo rapporto vengono applicate nuove metodologie di identificazione.

Per identificare gli eventi di intrusione stratosferica, in accordo con la metodologia già adottata presso la stazione globale GAW/WMO dello Zugspitze in Germania (si veda Trickl et al., 2010), vengono selezionati i periodi di misura caratterizzati da: **i) RH inferiore a 60%, (ii) valori minimi di RH inferiori a 30%; (iii) masse d'aria con vorticità potenziale superiore a 1.6 pvu.**

Per identificare la presenza di masse d'aria inquinate a CMN, si selezionano i periodi di misura per i quali i valori medi giornalieri di almeno due inquinanti "primari" (CO, NO, BC, NO₂) siano superiori alle loro medie mensili climatologiche.

Reference: Trickl, T., Feldmann, H., Kanter, H.-J., Scheel, H.-E., Sprenger, M., Stohl, A., and Wernli, H.: Forecasted deep stratospheric intrusions over Central Europe: case studies and climatologies, *Atmos. Chem. Phys.*, 10, 499-524, doi:10.5194/acp-10-499-2010, 2010.

List of parameters

In the following table, we provide the list of the atmospheric parameters observed at ICO-OV, together with a brief description of their key roles in the atmospheric investigations and the experimental set-up.

Parameters	Key role	Instrumentation
Surface ozone	Short-lived climate forcer, greenhouse gas, secondary pollutant	UV-absorption analyser (<i>Thermo Tei 49i</i>)
Carbon monoxide	Primary pollutant, ozone precursor, combustion tracer	Non dispersive IR absorption (<i>Thermo Tei48c-TL</i>)
Nitrogen oxides	Primary (NO) and secondary (NO ₂) pollutants, ozone precursors, combustion tracers.	Chemiluminescence analyser (<i>Thermo Tei42i-TL</i>)
Sulphur dioxide	Primary pollutant, secondary aerosol precursor, volcanic tracer.	UV-fluorescence analyser (<i>Thermo 43i-TLE</i>)
Black carbon	Short-lived climate forcer, primary pollutant, combustion tracer. It contributes to PM ₁	Multi-Angle Absorption Photometer (<i>Thermo MAAP 5012</i>)
Aerosol light scattering coefficient	Investigation of the aerosol climate direct forcing	3 –wavelenghts Integrating nephelometer (<i>TSI 3563</i>)
Aerosol number concentration (fine)	Short-lived climate forcer, primary and secondary aerosol, pollution tracer. It contributes to PM ₁ .	Optical particle counter (<i>GRIMM 1108</i>)
Aerosol number concentration (coarse)	Short-lived climate forcer, primary aerosol, mineral dust and sea salt tracer. It contributes to PM ₁₀ .	Optical particle counter (<i>GRIMM 1108</i>)
Halogenated gases	Stratospheric ozone depleting substances and climate forcer	Gas chromatography-Mass spectrometry. (<i>Agilent 6850–5975</i>)
Volatile organic compounds	Ozone and PM precursors	Gas chromatography-Mass spectrometry (<i>Agilent 6850–5975</i>)
Stratospheric nitrogen dioxide	Ozone destroying substance and buffer against halogen catalysed ozone loss	GASCOD-MTC: UV-Vis spectrometer
Temperature and relative humidity	Meteorology and data interpretation	<i>Rotronic, IRDAM WS 7000</i>
Atmospheric pressure	Meteorology and data interpretation	<i>Technoel, IRDAM WS 7000</i>
Wind	Meteorology and data interpretation	<i>Vaisala WS425, IRDAM WST7000</i>
Solar radiation	Meteorology and data interpretation	Silicon cell pyranometer (<i>Skye SKS110</i>)
UV-B radiation	Meteorology and data interpretation	Silicon photodiode (<i>Skye SKU 430</i>)

Lista dei parametri

Nella tabella è presentata la lista dei parametri osservati, assieme ad una breve descrizione dei ruoli nelle ricerche condotte ed il set up sperimentale utilizzato presso l'ICO-OV.

Parametri	Ruolo chiave clima – qualità dell'aria	Strumentazione
Ozono superficiale	Forzante climatico a breve tempo di vita, gas serra, inquinante secondario.	Analizzatore ad assorbimento UV (<i>Thermo Tei 49i</i>)
Monossido di carbonio	Inquinante primario, precursore dell'ozono, tracciante della combustione	Analizzatore ad assorbimento infrarosso (<i>Thermo Tei48c-TL</i>)
Ossidi d'azoto	Inquinanti primari (NO) e secondari (NO ₂), precursori dell'ozono, tracciante della combustione. In stratosfera NO ₂ influenza le concentrazioni di ozono.	Analizzatore a chemiluminescenza (<i>Thermo Tei42i-TL</i>)
Anidride solforosa	Inquinante primario, precursore dell'aerosol secondario, tracciante di eruzioni vulcaniche	Analizzatore a fluorescenza UV (<i>Thermo Tei43i-TLE</i>)
Black carbon	Forzante climatico a breve tempo di vita, inquinante primario, tracciante della combustione. Contribuisce al PM ₁	Fotometro per l'assorbimento multiangolare (<i>Thermo MAAP 5012</i>)
Coefficiente di scattering delle particelle	Studio della forzante climatica diretta degli aerosol	Nefelometro a 3 lunghezze d'onda(<i>TSI 3563</i>)
Numero delle particelle fini	Forzante climatico a breve tempo di vita, aerosol primario e secondario, tracciante dell'inquinamento. Contribuisce al PM ₁	Contatore ottico (<i>GRIMM 1108</i>)
Numero delle particelle grossolane	Forzante climatico a breve tempo di vita, aerosol primario, tracciante delle polveri minerali e del sale marino. Contribuisce al PM ₁₀	Contatore ottico (<i>GRIMM 1108</i>)
Gas alogenati	Distruggono l'ozono stratosferico, forzanti climatici	Gas cromatografia-Spettrometria di massa (<i>Agilent 6850–5975</i>)
Composti organici volatile	Precursori dell'ozono troposferico e del PM	Gas cromatografia-Spettrometria di massa (<i>Agilent 6850–5975</i>)
Biossido di azoto stratosferico	Distrugge l'ozono stratosferico e sostanza "tampone" per alogeni attivi nella deplezione dell'ozono stratosferico	Spettrometro UV/Vis GASCOD-MTC
Temperatura ed umidità relativa	Meteorologia ed interpretazione dei dati	<i>Rotronic, IRDAM WS 7000</i>
Pressione atmosferica	Meteorologia ed interpretazione dei dati	<i>Technoel, IRDAM WS 7000</i>
Vento	Meteorologia ed interpretazione dei dati	<i>Vaisala WS425, IRDAM WST7000</i>
Radiazione solare	Meteorologia ed interpretazione dei dati	Piranometro a celle di silicio (<i>Skye SKS110</i>)
Radiazione UV-B	Meteorologia ed interpretazione dei dati	Fotodiodo al silicio (<i>Skye SKU 430</i>)

Summary

OVERWIEV

Spring 2015 did not present high average levels of **short-lived climate forcers (SLCF)**: a value lower than the climatological mean was observed for **black carbon, surface ozone and fine particles**, while **carbon monoxide** and **coarse particles** average concentrations were well comparable with the typical spring values.

The 33% of the spring season was affected by **pollution events**, with the large number of events on March (14/31), especially from March, 11st to 27th.

26 days (27.9%) were affected by **mineral dust transport**: the event occurring from May 3rd to 8th showed the highest **coarse particle** average concentration.

The selection methodology allow the identification of 33 days (35.7 %) related with air-mass **transport from the stratosphere**. The period from March 12th to 17th was continuously affected, on a daily basis, by the occurrence of SI events.

Daily **surface ozone** peak was recorded on 12-05 (66.0 ppb). 30-minute average values ranged from a minimum of 34.1 ppb (03-04) to 76.2 ppb (11-05), with an average seasonal value of 55.7 ppb. This value is lower than the average climatological spring value obtained from the last 19 years (58.7 ppb).

Daily **carbon monoxide** peak was recorded on 17-03 (221.3 ppb). 30-minute average values ranged from a minimum of 80.2 ppb (08-05) to 263.7 ppb (17-03), with an average seasonal value of 135.1 ppb. This value is comparable to the average spring value obtained from the last 2 years (134.6 ppb).

Daily **nitric oxide** and **nitrogen dioxide** peaks were recorded on 12-03 (0.42 ppb) and 13-03 (1.05 ppb), respectively. 30-minute average values ranged from values below the detection limit to 3.22 ppb (for NO) and 3.35 ppb (for NO₂).

Daily **sulfur dioxide** peak was recorded on 18-03 (0.44 ppb). 30-minute average values ranged from values below the detection limit to 1.73 ppb (18-03), with an average seasonal value below the detection limit.

Daily **black carbon** peak was recorded on 20-03 (706.6 ng m⁻³). 30-minute average values ranged from a minimum of 10.0 ng m⁻³ (28-04) to 1260.3 ng m⁻³ (11-03), with an average seasonal value of 195.7 ng m⁻³. This value is lower than the average climatological spring value obtained from the last 10 years (229.5 ng m⁻³).

Sommario

VISIONE DI INSIEME

La primavera del 2015 non ha presentato valori medi elevati di *short-lived climate forcers* (SLCF): il **black carbon**, le **particelle fini** e l'**ozono** hanno mostrato valori inferiori rispetto alle medie climatologiche, mentre le concentrazioni di **monossido di carbonio** ed i valori delle **particelle grossolane** risultano essere in linea con quanto solitamente osservato durante la stagione primaverile.

Il 33 % della stagione è stato influenzato da eventi di **inquinamento**, con una prevalenza nel mese di marzo, quando sono stati osservati 14 dei 31 eventi.

Eventi di trasporto di sabbia sahariana hanno interessato il sito di misura per 26 giorni (27.9% della stagione).

La metodologia di selezione ha permesso di identificare 33 eventi associabili a trasporto di **masse d'aria provenienti dalla stratosfera**: su base giornaliera, il periodo compreso fra il 12 ed il 17 marzo è stato costantemente interessato da tali eventi.

Il valore massimo giornaliero della concentrazione di **ozono superficiale** è stato registrato il 12-05 (66.0 ppb). Le medie semi-orarie variano da 34.1 ppb (03-04) a 76.2 ppb (11-05), con un valore medio stagionale di 55.7 ppb. Tale valore è inferiore a quello climatologico relativo agli ultimi 19 anni (58.7 ppb).

Il valore massimo giornaliero della concentrazione di **monossido di carbonio** è stato registrato il 17-03 (221.3 ppb). Le medie semi-orarie variano da 80.2 ppb (08-05) a 263.7 ppb (17-03), con un valore medio stagionale pari a 135.1 ppb. Tale valore è confrontabile con quello delle ultime due stagioni primaverili (134.6 ppb).

I valori massimi giornalieri di **ossido d'azoto** e **biossido d'azoto** sono stati registrati rispettivamente il 12-03 (0.42 ppb) e il 13-03 (1.05 ppb). Le medie semi-orarie sono variate da valori inferiori al limite di rivelazione sino a 3.22 ppb (per NO) e 3.35 ppb (per NO₂).

Il valore massimo giornaliero di **biossido di zolfo** è stato registrato il 18-03 (0.44 ppb). Le medie semi-orarie sono variate da valori inferiori al limite di rilevabilità sino a 1.73 ppb (18-03), con un valore medio stagionale inferiore al limite di rilevabilità.

Il valore massimo giornaliero della concentrazione di **black carbon** è stato registrato il 20-03 (706.6 ng m⁻³). Le medie semi-orarie variano da 10.0 ng m⁻³ (28-04) a 1260.3 ng m⁻³ (11-03), con un valore medio stagionale pari a 195.7 ng m⁻³. Tale valore è inferiore a quello climatologico relativo agli ultimi 10 anni (229.5 ng m⁻³).

Daily **fine aerosol particles** peak was recorded on 20-03 (132.3 cm^{-3}). 30-minute average values ranged from below the detection limit to 261.7 cm^{-3} (11-03), with an average seasonal value of 20.1 cm^{-3} . This value is lower than the average climatological spring value obtained from the last 13 years (28.7 cm^{-3}).

Daily **coarse aerosol particles** peak was recorded on 05-05 (1.4 cm^{-3}). 30-minute average values ranged from below the detection limit to 2.1 cm^{-3} (05-05), with an average seasonal value of 0.2 cm^{-3} . This value is comparable to the average climatological spring value obtained from the last 13 years (0.3 cm^{-3}).

Daily **aerosol scattering coefficient at 550 nm** peak was recorded on 14-03 (58.8 Mm^{-1}). 30-minute average values ranged from below the detection limit to 150.2 Mm^{-1} (11-03), with an average seasonal value of 11.1 Mm^{-1} . This value is considerably lower than the average climatological summer value obtained from the last 8 years (26.2 Mm^{-1}).

Il valore massimo giornaliero della concentrazione di **particelle fini** è stato registrato il 20-03 (132.3 cm^{-3}). Le medie semi-orarie variano da valori inferiori al limite di rivelazione a 261.7 cm^{-3} (11-03), con un valore medio stagionale pari a 20.1 cm^{-3} . Tale valore è inferiore a quello climatologico degli ultimi 10 anni (28.7 cm^{-3}).

Il valore massimo giornaliero della concentrazione di **particelle grossolane** è stato registrato il 05-05 (1.4 cm^{-3}). Le medie semi-orarie variano da valori inferiori al limite di rivelazione a 2.1 cm^{-3} (05-05), con un valore medio stagionale pari a 0.2 cm^{-3} . Tale valore è in linea con quello climatologico relativo agli ultimi 13 anni (0.3 cm^{-3}).

Il picco giornaliero del **coefficiente di scattering dell'aerosol a 550 nm** è stato osservato il 14-03 (58.8 Mm^{-1}). Le medie sui 30-minuti oscillano tra valori inferiori al limite di rilevabilità e 150.2 Mm^{-1} (11-03), con un valore medio stagionale di 11.1 Mm^{-1} che risulta essere considerevolmente inferiore al valore medio climatologico relativo agli ultimi 8 anni (26.2 Mm^{-1}).

Daily **air temperature** peak was recorded on 05-05 (11.7°C), minimum on 07-03 (-9.3°C). 30-minute average values ranged from a minimum of -11.7°C (07-03) to 15.9°C (11-05), with an average seasonal value of 1.6°C , which is slightly higher than the seasonal climatological value (1.2°C).

Daily **relative humidity** minimum was recorded on 16-04 (33.4%). 30-minute average values ranged from a minimum of 9.2 % (20-04) to a maximum of 100.0 % (observed on 25 days), with an average seasonal value of 79.2 %. This value is comparable to the average climatological spring value obtained from the last 19 years (81.4 %).

Daily **atmospheric pressure** peak was recorded on 13-04 (800.1 hPa), the lowest value on 25-03 (773.6 hPa). 30-minute average values ranged from a minimum of 769.0 hPa (25-03) to 801.1 hPa (12-04), with an average seasonal value of 789.8 hPa, which is slightly higher than the average climatological spring value obtained from the last 19 years (787.4 hPa).

Daily **wind speed** peak was recorded on 07-03 (25.2 m s^{-1}). 30-minute average values ranged from a minimum of 0.7 m s^{-1} (14-03) to a maximum of 36.9 m s^{-1} (08-03), with an average seasonal value of 6.8 m s^{-1} . This value is slightly lower than the climatological spring value (7.9 m s^{-1}).

Wind direction was prevalently SW (34.1 % of 30-minute data), in line with the climatological analysis over the last 19 years .

Daily **solar radiation** highest average daily value was recorded on 29-05 (368.3 W m^{-2}), with clear sky conditions observed throughout. The lowest average daily value (13.6 W m^{-2}) was observed on 01-03, during a period (from 01-03 to 07-03) when cloud cover was present at the measurement site. Snow and ice over sensors could have affected measurements.

A similar trend was also observed for **UV-B radiation**, with the highest value observed on 29-05 (0.60 W m^{-2}). Snow and ice over sensors could have affected measurements.

Il valore massimo giornaliero della **temperatura** è stato registrato il 05-05 (11.7 °C), il valore minimo il 07-03 (-9.3 °C). Le medie semi-orarie variano da -11.7 °C (07-03) a 15.9 °C (11-05), con un valore medio stagionale pari a 1.6 °C, leggermente superiore a quello medio climatologico primaverile (1.2 °C).

Il valore minimo giornaliero dell'**umidità relativa** è stato registrato il 16-04 (33.4 %). Le medie semi-orarie variano da 9.2 % (20-04) a 100 % (osservato in 25 giornate), con un valore medio stagionale pari a 79.2 %. Tale valore è confrontabile con quello climatologico relativo agli ultimi 19 anni (81.4 %).

Il valore massimo giornaliero della **pressione atmosferica** è stato registrato il 13-04 (800.1 hPa), il valore minimo il 25-03 (773.6 hPa). Le medie semi-orarie variano da 769.0 hPa (25-03) a 801.1 hPa (12-04), con un valore medio stagionale pari a 789.8 hPa, leggermente superiore a quello climatologico relativo agli ultimi 19 anni (787.4 hPa).

Il valore massimo giornaliero della **velocità del vento** è stato registrato il 07-03 (25.2 m s⁻¹). Le medie semi-orarie variano da 0.7 m s⁻¹ (14-03) a 36.9 m s⁻¹ (08-03), con un valore medio stagionale pari a 6.8 m s⁻¹. Tale valore è leggermente inferiore rispetto a quello climatologico ottenuto dalle misure realizzate negli ultimi 19 anni (7.9 m s⁻¹).

La **direzione del vento** osservata nell' inverno 2015 è stata prevalentemente da Sud-Ovest (34.1 % dei dati semi-orari), in linea con la climatologica prevalenza di venti sudoccidentali.

La **radiazione solare** mostra il valore giornaliero massimo il 29-05 (368.3 W m⁻²), con valori elevati associati a condizioni di cielo libero da nubi osservate per quasi tutta la stagione. Il minimo giornaliero (13.6 W m⁻²) è stato osservato il 01-03, all'interno di un periodo compreso tra il 01-03 ed il 07-03 caratterizzato dalla presenza di copertura nuvolosa. Possibili accumuli di neve e ghiaccio possono inoltre aver indotto l'occorrenza di bassi valori di radiazione.


Analogo comportamento viene osservato per quanto riguarda la **radiazione UV**, con il massimo giornaliero, pari a 0.60 W m⁻², osservato il 29-05. Possibili accumuli di neve e ghiaccio possono inoltre indurre l'occorrenza di bassi valori di radiazione UVB.

Special events

In this paragraph, we present a detailed overview of “special events” which have been detected at the ICO-OV during the reference period, namely:

- Mineral Dust transport
- Stratospheric intrusions;
- Pollution transport;

Please note that the event selection methodologies are executed on 30-minute basis, thus, for the same day, different classes of special events can be observed.



LEGEND






Eventi speciali

In questo paragrafo viene presentato l'elenco degli "eventi speciali" che sono stati registrati presso il sito di misura durante il periodo analizzato:

- **Trasporto di polveri minerali;**
- **Intrusioni stratosferiche;**
- Trasporto di inquinanti

Va notato che le metodologie di selezione degli eventi sono calcolate a partire dai dati a 30-minuti, quindi, per lo stesso giorno, possono essere osservate diverse tipologie di evento.



LEGENDA

Trasporto polveri minerali Intrusioni stratosferiche Trasporto di inquinanti


Mineral dust transport

The presence of mineral dust in the atmosphere plays direct and indirect role in affecting climate variations. Moreover, mineral dust can seriously affect air quality in regions downwind of desert areas, contributing to PM₁₀ levels. Sahara desert exports more mineral dust than any other area of the world, injecting into the atmosphere millions of Tons of dust particles. Mt. Cimone represents one of the first mountain ridges that Saharan dust meet along their tracks towards Italy and Europe.

Selection methodology: we detected a Saharan dust event when the atmospheric concentration of coarse particles (particles with diameter $1 \mu\text{m} \leq D_p \leq 20 \mu\text{m}$) significantly increased with air-masses coming from North Africa, as deduced by three-dimensional air-mass back-trajectories and transport model outputs.

WINTER 2015:

- 26 days were characterized by the transport of mineral dust from northern Africa (27.9 % of the period).
- The mineral dust transport event are evenly distribute throughout the season, with six events characterized by a long duration.
- During the dust event of 5th May the daily average coarse particle number concentration peaked at 1.4 cm^{-3} .



Dust transport event simulation by NAAPS model (5th May 2015).

Simulazione dell'evento di trasporto di polveri minerali osservato il 05 Maggio 2015 (modello NAAPS).

<http://www.nrlmry.navy.mil/>

Courtesy by NRL/Monterey Aerosol Modeling.

Trasporto di polveri minerali

La presenza di aerosol (polveri) minerali nell'atmosfera può influenzare il clima attraverso effetti diretti ed indiretti. Esse possono inoltre alterare in modo significativo la qualità dell'aria in regioni prossime alle aree di emissione o soggette a fenomeni di trasporto, influenzando le concentrazioni di PM₁₀. Masse d'aria ricche di polveri minerali possono essere trasportate dal deserto del Sahara, la più importante sorgente al mondo di polveri minerali, verso l'Italia e l'Europa. Mt. Cimone rappresenta uno dei primi rilievi montuosi che queste masse d'aria incontrano durante il loro movimento verso nord.

Metodologia di selezione: gli eventi di trasporto di polveri sahariane sono stati identificati quando la concentrazione delle particelle grossolane ($1 \mu\text{m} \leq D_p \leq 20 \mu\text{m}$) è aumentata in modo significativo con l'arrivo di masse d'aria provenienti dal nord Africa, come indicato da analisi di retro-traiettorie tri-dimensionali delle masse d'aria e da modelli di trasporto.

INVERNO 2015:

- **26 giorni sono stati caratterizzati dal trasporto di polveri minerali proveniente dal Nord Africa (27.9 % del periodo).**
- **Gli eventi di trasporto sono uniformemente distribuiti nei 3 mesi, con 6 eventi principali caratterizzati da elevata durata temporale.**
- **Il picco di concentrazione di particelle grossolane (1.4 cm^{-3}) è stato osservato il 5 maggio 2015.**



Dust transport observed by MODIS sensor on-board on AQUA and TERRA NASA satellite (5th May 2015). The brownish clouds represent mineral dust transported toward Monte Cimone.

Rappresentazione tramite immagine a colori dei satelliti Terra/Aqua dell'evento di trasporto di polveri minerali osservato il 05 Maggio 2015. Sono evidenti le masse d'aria ricche di aerosol minerale che vengono trasportate verso Monte Cimone.

<https://earthdata.nasa.gov>

Courtesy by NASA - Worldview

Stratospheric intrusions (SI)


Stratospheric intrusions (SI) can be considered as a specific aspect of stratosphere–troposphere exchange (STE): the irreversible downward transport of stratospheric air relatively deep into the troposphere. Such phenomena are highly episodic and can be favored by a number of different mechanisms, acting on different geographical and temporal scales: tropopause folding and cut-off lows at upper levels, and fronts or high-pressure systems at the surface.

Even though it has been assessed that nowadays the greatest contribution to tropospheric ozone concentrations comes from photochemical production, the contribution from STE is far from negligible, in particular in the free troposphere. For these reasons, at ICO-OV the frequency of SI and its contribution to ozone is assessed.

Selection methodology: at Mt. Cimone, we identified days probably affected by air-mass transport from the stratosphere or from the upper free troposphere by selecting the measurement periods characterized by at least 6 hours of relatively dry conditions ($\text{RH} < 60\%$) during which RH also reached a value below 30 %, together with analysis of air-mass three-dimensional back-trajectories corroborating the origin of the air masses (daily potential vorticity maximum at least 1.6 ppu). **In respect to the former reports, this methodology has been updated in agreement with Trickl et al. (2010).**

WINTER 2015:

- During spring 2015, we detected 33 days affected by the transport of air masses from the stratosphere, representing the 35.7 % of the analysed period.
- The highest ozone concentration during a STE was observed on April 12th, with a daily average of 65.7 ppb.
- March 12th – 17th represents the longest seasonal period continuously interested by the presence of Stratospheric Intrusions.



Trajectories describing the path of stratospheric air-masses for the event of 12 April 2015. The color code represents the air-mass height (expressed as pressure level).

Courtesy by Michael Sprenger (ETH-Z, Switzerland)

Traiettorie che descrivono il moto in atmosfera di masse d'aria d'origine stratosferica per l'evento del 12 aprile 2015. Il colore rappresenta la quota (espressa come livello di pressione) delle masse d'aria.

Elaborazione: Michael Sprenger (ETH-Z, Switzerland)


Intrusioni stratosferiche (SI)

Le intrusioni stratosferiche (SI) possono essere considerate un aspetto specifico degli scambi stratosfera-troposfera (STE). Tali fenomeni, che avvengono in maniera episodica, possono essere favoriti da processi dinamici e meteorologici caratteristici che agiscono su differenti scale spazio-temporali: ripiegamento della tropopausa, cut-off low, sistemi frontali o aree di alta pressione. Sebbene il processo più importante che influenza la variabilità dell'ozono in troposfera sia oggi rappresentato dalla produzione fotochimica, il contributo dei processi STE è tutt'altro che trascurabile, in particolare nella libera troposfera. Per queste ragioni, presso l'ICO-OV viene effettuata l'identificazione e lo studio di questa classe di fenomeni.

Metodologia di selezione: a Mt. Cimone, sono stati identificati gli eventi di trasporto di masse d'aria dalla stratosfera o dalla parte superiore della libera troposfera come i periodi caratterizzati per almeno 6 ore dalla presenza di masse d'aria relativamente secche ($\text{RH} < 60\%$) e che abbiano presentato valori minimi di RH inferiori al 30%. Retro-traiettorie tridimensionali delle masse d'aria, sono state utilizzate per corroborare l'origine degli eventi (valore massimo giornaliero della vorticità potenziale superiore a 1.6 pvu). **La metodologia di selezione è stata modificata rispetto all'ultimo rapporto stagionale, in accordo con la procedura presentata da Trickl et al. (2010).**

INVERNO 2015:

- Durante l'inverno 2015 sono stati identificati 33 episodi di intrusione stratosferica, corrispondenti al 35.7 % del periodo analizzato.
- Il 12 Aprile è stata osservata la concentrazione stagionale di ozono più elevata durante un evento di intrusione stratosferica (valore medio giornaliero: 65.7 ppb).
- Il periodo compreso tra il 12 ed il 17 Marzo è stato caratterizzato dalla continua presenza di intrusioni stratosferiche.




Pollution transport

The Mediterranean region represents a global hot-spot in terms of climate change and atmospheric composition variability while the Po Basin on which Mt. Cimone leans out, is considered one of the major polluted European regions. In particular during the summer seasons, when the high solar irradiance characterized these areas, many anthropogenic pollutants, including photochemically produced ozone can affect the lower troposphere. With the goal of better evaluating the influence of these processes on the atmospheric composition variability, polluted air-mass transport phenomena are systematically identified and investigated at ICO-OV.

Selection methodology: at Mt. Cimone, days possibly affected by polluted air-mass transport are identified by selecting the measurement periods for which at least two “primary” pollutants (CO, NO, BC, NO₂) showed average daily values higher than the monthly climatological mean value. **This methodology has been updated with regard to the former reports, with the aim of provide a more robust and smart event detection.**

WINTER 2015

- 31 days, were characterized by significant transport of polluted air masses (33 % of the period).
- March was the month the most affected by pollution events (14 days), with 13 of them continuously affecting CMN from March 11th until 24th.



Open fire emission transport simulation by NAAPS model (20th March 2015). The colored scale represents the smoke aerosol concentration expressed as $\mu\text{g m}^{-3}$.

Simulazione del trasporto di emissiozioni da incendi il 20 marzo 2015 (modello NAAPS). La scala colorata rappresenta la concentrazione dell'aerosol emesso dagli incendi in $\mu\text{g m}^{-3}$.

<http://www.nrlmry.navy.mil/>

Courtesy by NRL/Monterey Aerosol Modeling


Trasporto di inquinanti

Il bacino del Mediterraneo rappresenta un “hot-spot” globale per quanto riguarda i cambiamenti del clima e della composizione dell’atmosfera, mentre la Pianura Padana rappresenta un’importante area sorgente di inquinamento antropico. In particolare durante l'estate, a causa dell'alto irraggiamento solare che caratterizza queste regioni, esse sono interessate da intensi eventi di produzione fotochimica e di ozono nella bassa troposfera. Con lo scopo di valutare l'influenza di tali eventi sulla composizione dell'atmosfera, i fenomeni di trasporto di masse d'aria inquinate sono sistematicamente identificati e studiati a Mt. Cimone

Metodologia di selezione: a Mt. Cimone, sono stati identificati i giorni possibilmente affetti da trasporto di masse d'aria inquinate selezionando i periodi di misura per i quali i valori medi giornalieri di almeno due inquinanti “primari” (CO, NO, BC, NO₂) siano superiori alle loro medie mensili climatologiche. **La metodologia è modificata rispetto ai precedenti rapporti stagionali, al fine di rendere l'identificazione degli eventi più semplice e robusta.**

AUTUNNO 2014

- 31 giorni (33% della stagione), sono stati caratterizzati da episodi significativi di trasporto di masse d'aria inquinate.
- Durante il mese di marzo, si sono verificati 14 dei 31 episodi di trasporto di inquinanti individuati: il periodo 11-25 marzo è stato costantemente interessato da questa classe di eventi.



Surface ozone

Why is ozone so important?

Ozone (O_3) is one of the most important Short-Lived Climate Forcers and Pollutant (SLCF/P), being a powerful greenhouse gas at regional scale. Due to its chemical properties, O_3 is also a dangerous secondary pollutant in the lower troposphere. Its tropospheric mixing ratios are also affected by natural processes, e.g. stratospheric intrusions and lightning production. Being the precursor of oxidizing substances like OH radical and NO_3 , O_3 is one of the key agents determining the oxidation capacity of the troposphere.

Instrumentation and calibration

Surface ozone is measured by using a UV-absorption analyser (Thermo Tei 49i). Intercomparisons with the laboratory standard (Thermo 49iPS, traced back to SRP#15 at the World Calibration Centre for surface ozone at WCC-EMPA of Zürich) are carried out every 3-months.


Basic statistical parameters

Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (%)	Min value (ppb)	25 th Percentile (ppb)	50 th Percentile (ppb)	Average mean value (ppb)	75 th percentile (ppb)	Max value (ppb)
93.2	34.1	51.0	56.1	55.7	60.5	76.2


Time series of daily mean values

As expected general increase in ozone concentration is observed throughout the season, representing the natural transition from the winter minimum to the spring/summer maximum. High concentrations were observed in mid-April and May: on May 12th the highest daily average O_3 was observed (66.0 ppb) during a pollution transport event.



Comparison with historical data-set

The O₃ seasonal mean value is 55.7 ppb, a value **lower than the climatological one (58.7 ppb)**. This was related to the lower than average concentrations during the first half of the season, partially compensated by higher concentration during mid-April and late May, when many events of stratospheric intrusions and pollution transport were observed at CMN. **It should be noted that in the last 7 years the inter-annual variability of springtime O₃ concentration is lower than in the previous decade.**



Carbon monoxide (NDIR)

Why is carbon monoxide so important?

Carbon Monoxide (CO) plays an important role in the oxidation/reduction chemistry of the atmosphere and it participates in the reactions of photochemical O₃ production. CO has an indirect radiative forcing effect by influencing atmospheric mixing ratios of O₃ and methane. Through natural processes in the atmosphere, CO is eventually oxidized to CO₂. CO represents a tracer for combustion emissions (biomass burning, residential, traffic,...).

Instrumentation and calibration

Carbon monoxide is measured by using a non-Dispersive Infrared (NDIR) analyzer (Thermo Scientific TEI 48C-TL). A CO working standard (approx. 10 ppm, synthetic air, Messer Italia) is used to calibrate the instrument with a dilution system. On a monthly basis, these working standards were compared against secondary standards from NOAA-CMDL.


Basic statistical parameters

Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (%)	Min value (ppb)	25 th Percentile (ppb)	50 th Percentile (ppb)	Average mean value (ppb)	75 th percentile (ppb)	Max value (ppb)
97.6	80.2	119.9	129.6	135.1	144.1	263.7


Time series of daily mean values

During march, CO was characterized by a strong variability with daily average concentration as low as 96.3 ppb on March 6th (stratospheric intrusion), followed by the highest mixing ratio (221.3 ppb) on March 17th during an episode of pollution transport. March appeared as the most "polluted" month, with the highest CO values and the highest occurrence of pollution events.



Comparison with historical data-set

The 2015 spring average mean value of CO was 135.1 ppb, which is **comparable to the average mean value of 134.6 ppb obtained from the last two Spring seasons.**



Nitrogen oxides

Why are nitrogen oxides so important?

Nitrogen oxides (NO_x) encompasses nitric oxide (NO) and nitrogen dioxide (NO_2). NO is naturally produced by lightning. Anthropogenic contributions are related to combustion processes and agricultural fertilization. NO_x are key elements of atmosphere chemistry influencing a number of atmospheric compounds with roles on climate, air-quality and ecosystem threats, e.g. sulphur dioxide, halocarbons, methane, tropospheric ozone, secondary aerosols.

Instrumentation and calibration

Nitrogen oxides ($\text{NO}_x = \text{NO} + \text{NO}_2$) are measured by using a Chemiluminescence analyser (Thermo 42i-TL), equipped with a photolytic converter (Blue Light Converter) for NO_2 determination. Every 48 hours, zero and span checks are carried out for NO by using an external zero air source (dry compressed air scrubbed with active charcoal and Purafill) and dilution of certified NO standard (5 ppm +/- 2%). GPT is used to determine the conversion efficiency of the NO_2 converter.

Basic statistical parameters


Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (%)	Min value (ppb)	25 th percentile (ppb)	50 th percentile (ppb)	Average mean value (ppb)	75 th percentile (ppb)	Max value (ppb)
NO 97.7	UDL	0.01	0.03	0.1	0.08	3.2
NO_2 97.7	UDL	0.07	0.13	0.2	0.26	3.4

UDL: under detection limit

Time series of daily mean values



The highest NO value (blue line: 0.42 ppb) was observed on March 12th, while the highest NO_2 value (red line: 1.05 ppb) was observed on March 13th. The highest NO_x values were observed during March, when the highest occurrence of pollution events were observed at CMN.



Comparison with historical data-set

The 2015 spring average mean value of NO (NO_2) was 0.07 ppb (0.22 ppb) which is **comparable (lower)** to the average spring mean value of **0.08 ppb (0.86 ppb)**.

It should be noted that NO_2 data were obtained by analyzer equipped by Molybdenum during seasons 2012 - 2013. These data can be significantly overestimated in respect to the observations made by photolytic converter (year 2013) due to the interference of processed N-oxidised species (i.e. PAN) to the former data series.



Sulphur dioxide

Why is sulfur dioxide so important?

Sulfur dioxide (SO_2) is the main precursor to the sulphate aerosol which exerts a large influence on climate and air quality. It is a well known precursor for acid rains. SO_2 is also one of the main tracers for detecting the occurrence of ash plumes from volcanic eruption. In Europe, during the last years, SO_2 concentrations have been declining due to efficient restrictions on emissions.

Instrumentation and calibration

Sulphur dioxide is measured by using a UV-fluorescence analyser (Thermo 43i-TLE). Daily zero check are executed by using an external zero air source (scrubber with active charcoal), while daily span check are performed using a permeation tube with set point at 48 ppb. Detection limit for 1-minute average is estimated to be 0.11 ppb

Basic statistical parameters


Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (%)	Min value (ppb)	25 th percentile (ppb)	50 th percentile (ppb)	Average mean value (ppb)	75 th percentile (ppb)	Max value (ppb)
98.0	UDL	UDL	UDL	UDL	0.16	1.7

UDL: under detection limit

Time series of daily mean values

During spring 2015 the average SO_2 mixing ratio was lower than the detection limit, even though several SO_2 peaks were observed throughout the season, especially on March (together with enhanced NO_x and CO values) and from May, 5th till 20th. The highest daily value (0.44 ppb) was observed on March 18th, during a pollution events.



NOTE: Negative values in the graph represent “under detection limit” conditions, common in a remote, high altitude, free troposphere measurement site.

Black carbon

Why is black carbon so important?

Black carbon (BC) is a primary aerosol resulting from incomplete combustion processes. Its main sources are fossil fuel combustion (anthropogenic) and biomass burning (natural and anthropogenic). BC, a Short Lived Climate Forcer and Pollutant, strongly absorbs solar radiation and it has been recognized as a driving factor of global warming: the magnitude of the direct radiative forcing due to BC can exceed that due to methane.

Instrumentation and calibration

Equivalent black carbon concentration is measured by a Multi Angle Absorption Photometer (MAAP, Model 5012 – Thermo Electron Corporation). Detection limit was measured as 3σ of 12 h measurement of free particle air. Calibration of sampling flow and internal temperature-pressure sensors are conducted every 6 months.


Basic statistical parameters

Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (%)	Min value (ng m ⁻³)	25 th percentile (ng m ⁻³)	50 th percentile (ng m ⁻³)	Average mean value (ng m ⁻³)	75 th percentile (ng m ⁻³)	Max value (ng m ⁻³)
95.3	10.0	78.8	143.8	195.7	246.1	1260.3


Time series of daily mean values

Similarly to what have been observed for other primary pollutants (CO, NO, SO₂) the highest BC average concentrations were observed on March and May. The seasonal daily average maximum (706.6 ng m⁻³) was observed on March 20th, with similarly high concentrations been observed on March 14th (630.5 ng m⁻³) during two pollution transport events.



Comparison with historical data-set

The 2015 spring average mean value of BC is 195.7 ng m^{-3} , which is lower than the climatological mean value (229.5 ng m^{-3}). This behavior is the result of lower than average concentrations throughout the season (only 30% of the day showed an average BC concentration higher than the climatological value), even though higher daily concentrations were observed during March.



Aerosol light scattering coefficient

Why is aerosol light scattering coefficient so important?

Aerosol light scattering coefficient variability is an important parameter in deriving quantitative information on the optical properties of atmospheric aerosols, which are used to determine the direct effects of aerosols on the earth radiation balance (and therefore their impact on climate change). Moreover, information on the extinction coefficient, which is related to visibility in the atmosphere, is also an important parameter in many atmospheric applications.

Instrumentation and calibration

Starting from March 2014 a new three wavelength TSI Integrating nephelometer 3563 measures the aerosol light scattering coefficient at red (700 nm), green (550 nm), and blue (450 nm) wavelengths. A calibration on site with low span gas (filtered air) and high span gas (filtered carbon dioxide) is performed every 3 months.

Basic statistical parameters


Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (%)	Min value (Mm^{-1})	25 th percentile (Mm^{-1})	50 th percentile (Mm^{-1})	Average mean value (Mm^{-1})	75 th percentile (Mm^{-1})	Max value (Mm^{-1})
700 nm 99.4	UDL	0.3	7.6	15.9	20.2	189.1
550 nm 99.4	UDL	0.2	5.1	11.1	13.6	150.2
450 nm 99.4	UDL	0.0	3.0	6.7	8.2	104.8

UDL: under detection limit

Time series of daily mean values


The highest daily mean value of the scattering coefficient at all three wavelength (450, 550, 700 nm) has been observed on March 14th, during the identified dust episode (respectively 82.2, 58.8 and $36.1 Mm^{-1}$). High values were observed (similarly to BC and SO₂) during March.



Comparison with historical data-set

The 2015 spring average mean value of scattering coefficient at 550 nm is 11.1 Mm^{-1} , which is lower than the climatological mean value (26.2 Mm^{-1}).

It should be noted that aerosol scattering data were obtained by a M9003 integrating nephelometer (ECOTECH) during autumn seasons 2007 - 2013.



Aerosol number concentration (fine)

Why are fine particle so important?

Fine particles are highly effective in modifying the radiation field by absorbing and scattering solar and thermal radiation, thus impacting radiative transfer through the atmosphere. Additionally, aerosols act as cloud condensation and ice nuclei, thus influencing cloud properties. Aerosols also help to control the concentrations, lifetime and the physical as well as the chemical behavior of many important trace gases by providing reaction sites and serving as carrier and/or sink for many atmospheric species. Moreover, fine particles strongly contribute to air pollution, representing a main fraction of PM₁.

Instrumentation and calibration

Aerosol concentration and size distribution of particles with optical diameter between 0.3 and 20 µm have been continuously recorded in 15-size channel by using an OPC Mod. GRIMM 1.108. These measurements allow the continuous measurement of the fine mode ($0.3 \mu\text{m} \leq D_p \leq 1 \mu\text{m}$) particle number. The instrument is based on the quantification of the 90° scattering of light by aerosol particles

Basic statistical parameters


Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability %	Min value (cm ⁻³)	25 th percentile (cm ⁻³)	50 th percentile (cm ⁻³)	Average mean value (cm ⁻³)	75 th percentile (cm ⁻³)	Max value (cm ⁻³)
94.5	0.0	4.5	10.1	20.1	21.2	261.7

UDL: under detection limit


Time series of daily mean values

Fine particles daily mean values were generally quite low throughout the season. Only on March high values were observed during pollution events. A high value was also observed on 12th, April when a transport of polluted air masses was further observed.



Comparison with historical data-set

Spring 2015 fine particles average number concentration was 20.1 cm^{-3} , **lower than the seasonal climatological value (28.7 cm^{-3})**. This behavior is similar to what is observed for other pollution tracers and is the result of clean conditions throughout the season, with the exception of March.



Aerosol number concentration (coarse)

Why is this research so important?

Coarse particles measured in background conditions represent a good tracer for mineral dust or marine aerosol transport. They play a significant role in radiation budget by absorbing and especially scattering solar radiation and can act as condensation and ice nuclei. Coarse particles can represent one of the major contributors to the overall PM₁₀ variability. Moreover, mineral dust contributes in determining the chemical behavior of many important trace gases (e.g. ozone) by way of heterogeneous-phase chemistry. Coarse particles strongly influence PM₁₀ concentrations.

Instrumentation and calibration

Aerosol concentration and size distribution of particles with optical diameter between 0.3 and 20 µm have been continuously measured in 15-size channel by using an OPC Mod. GRIMM 1.108. These measurements permit the determination of the coarse ($1 \mu\text{m} \leq D_p \leq 20 \mu\text{m}$) particle number. The instrument is based on the quantification of the 90° scattering of light by aerosol particles.

Basic statistical parameters


Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability %	Min value (cm ⁻³)	25 th percentile (cm ⁻³)	50 th percentile (cm ⁻³)	Average mean value (cm ⁻³)	75 th percentile (cm ⁻³)	Max value (cm ⁻³)
93.8	UDL	0.06	0.14	0.2	0.27	2.1

UDL: Under Detection Limit


Time series of daily mean values

With the exception of the high values observed during May that were related to dust transport events (highest daily average concentration of 1.4 cm⁻³ and 1.3 cm⁻³ were observed on May 5th and 14th), low coarse particles concentrations were observed throughout the season.



Comparison with historical data-set

The spring 2015 average mean value of the coarse particles (0.2 cm^{-3}) is **comparable to the climatological value (0.3 cm^{-3})**.



Air Temperature

Why is air-temperature so important?

Temperature data are useful to detect the occurrence of summer heat waves, during which photochemical smog episodes and transport of pollution from the boundary layer to the free troposphere can be favoured. The measurement of meteorological parameters at ICO-OV is a fundamental activity for the analysis of other measurements such as trace gases and aerosols.

Instrumentation and calibration

The basic meteorological data (temperature, relative humidity and atmospheric pressure) are measured above the ICO-OV terrace using instrumentation in compliance with WMO recommendations (IRDAM WST7000 and Rotronics thermo-hygrometer).


Basic statistical parameters

Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability %	Min value (°C)	25 th percentile (°C)	50 th percentile (°C)	Average mean value (°C)	75 th percentile (°C)	Max value (°C)
97.6	-11.7	-1.6	1.1	1.6	5.0	15.9


Time series of daily mean values

The air-temperature time series showed a clear increasing trend associated with the transition from winter to spring season. The highest daily mean value (11.7 °C) has been observed on May 5th, along with the highest coarse particle daily average number concentration, associated with a dust event.



Comparison with historical data-set

The spring 2015 average temperature (1.6°C) was **higher than the seasonal climatological value (1.2°C)**. This is the result of a stable weather regime characterizing the majority of the season, resulting in the measurement site been interested by warmer temperature and clear sky conditions.



Relative humidity

Why is relative humidity so important?

Relative humidity is a key parameter to identify the occurrence of dry meteorological conditions ($RH < 60\%$), usually associated with stratospheric intrusions or air-mass transport from the free troposphere. During summer, afternoon-evening RH increases can trace transport of air-masses from the boundary layer.

Instrumentation and calibration

The basic meteorological data (temperature, relative humidity and atmospheric pressure) are measured above the ICO-OV terrace using instrumentation in compliance with WMO recommendations (IRDAM WST7000 and Rotronics thermo-hygrometer).


Basic statistical parameters

Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (%)	Min value (%)	25 th percentile (%)	50 th percentile (%)	Average mean value (%)	75 th percentile (%)	Max value (%)
97.6	9.2	62.1	87.4	79.2	99.0	100.0


Time series of daily mean values

The daily mean RH values ranged from 100% to 9.2%. The 32.6% of days showed average value higher than 95%. Periods characterized by relatively dry conditions (mean daily values lower than 60%) were observed through the 8th - 22th April when significant occurrence of stratospheric intrusions were detected at the measurement site.



Comparison with historical data-set

The spring 2015 average relative humidity (79.2%) was comparable to the seasonal climatological value (81.4%).



Atmospheric pressure

Why is atmospheric pressure so important?

Pressure is a key parameter to investigate the variability of weather conditions at the ICO-OV. As an example, heat waves periods are characterized by the occurrence of high pressure values, while sudden pressure variability can be used to identify the passage of synoptic-scale disturbances possibly related to stratospheric intrusions.

Instrumentation and calibration

The basic meteorological data (temperature, relative humidity and atmospheric pressure) are measured above the ICO-OV terrace using instrumentation in compliance with WMO recommendations (IRDAM WST7000 and Tecnoel barometer).


Basic statistical parameters

Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (hPa)	Min value (hPa)	25 th percentile (hPa)	50 th percentile (hPa)	Average mean value (hPa)	75 th percentile (hPa)	Max value (hPa)
97.6	769.0	786.6	790.0	789.8	793.1	801.1


Time series of daily mean values

High pressure values were observed during the central part of April, when the maximum seasonal daily average value was recorded (800.1 hPa), together with the lowest relative humidity value, probably related to the presence of an anticyclonic dome which favored downward motion of air-masses in the lower troposphere.



Comparison with historical data-set

The spring 2015 averaged atmospheric pressure (789.8hPa) was **higher than the climatological value (787.4 hPa)**.



Wind speed and direction

Why is wind so important?

Wind speed and direction are used to identify the air mass circulation and therefore the transport of polluted air-masses from the near Po basin, as well as to identify the passage of surface fronts and the development of thermal wind circulation.

Instrumentation and calibration

Wind measurements are carried out at 5 m and 3 m high above the roof of the station, by using an integrated weather station IRDAM WST7000 and a sonic anemometer Vaisala WS425, respectively.



Basic statistical parameters of wind speed

Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (%)	Min value (m/s)	25 th Percentile (m/s)	50 th Percentile (m/s)	Average mean value (m/s)	75 th percentile (m/s)	Max value (m/s)
85.7	0.7	3.1	5.3	6.8	8.8	36.9



Time series of daily mean values

The observed prevailing directions shows a prevalence of SW wind, with wind speed being considerably higher during the first part of the season, while low values were observed from April 10th onward.



Comparison with historical data-set

The spring 2015 showed an average wind speed (6.8 m/s) lower than the climatological value (7.9 m/s). The seasonal wind prevalent direction is SW (34.1 %), followed by NE(17.5 %), in agreement with the prevailing wind direction observed at CMN.



Solar radiation (short-wave and UV-B)

Why is solar radiation so important?

Solar radiation is a key parameter in studying climate change and also play a role in defining the chemical properties of the troposphere, triggering photochemical reactions of important compounds (like O₃). Moreover, UV-B radiation is fundamental in determining the oxidative properties of the troposphere by leading O₃ photo-dissociation and thus determining OH levels.

Instrumentation and calibration

Solar radiation (wavelength: 350 – 1100 nm) and UV-B (wavelengths: 280–315 nm) are respectively measured by a commercial silicon cell pyranometer (Skye SKS110) and a silicon photodiode (Skye SKU 430). Calibrations were performed by factory against a WMO secondary standard pyranometer (for Skye SKS110) and against the National Physical Laboratory UK reference standard lamp (for Skye SKU 430).


Basic statistical parameters (Solar radiation)

Statistical parameters are calculated basing on 30-minute aggregated values from March 2015 to May 2015.

Data availability (%)	Min value (W/m ²)	25 th Percentile (W/m ²)	50 th Percentile (W/m ²)	Average mean value (W/m ²)	75 th percentile (W/m ²)	Max value (W/m ²)
95.6	UDL	UDL	21.1	179.0	287.0	1019.2

UDL: under detection limit

Time series (Solar radiation)




Basic statistical parameters (UV-B)

Statistical parameters are calculated basing on 30-minute aggregated values from December 2014 to February 2015.

Data availability (%)	Min value (W/m^2)	25^{th} Percentile (W/m^2)	50^{th} Percentile (W/m^2)	Average mean value (W/m^2)	75^{th} percentile (W/m^2)	Max value (W/m^2)
85.1	UDL	UDL	UDL	0.3	0.43	2.0

UDL: under detection limit

Time series (UV-B)



Bibliography

Here we present a list of the main scientific articles, from the year 2000 onward, resulted from the research activity conducted at ICO-OV:

Bonasoni P, Stohl A, Cristofanelli P, Calzolari F, Colombo T, Evangelisti F: Background ozone variations at Mt. Cimone, Atmos. Environ., 34 (29-30), 2000.

<http://www.sciencedirect.com/science/article/pii/S1352231000002685>

Wotawa G, Kroger H, Stohl A: Transport of ozone towards the Alps: results from trajectory analyses and photochemical model studies, Atmos. Environ., 34 (9), 2000.

<http://www.sciencedirect.com/science/article/pii/S1352231099003635>

Bonasoni P, Evangelisti F, Bonafè U, Ravegnani F, Calzolari F, Stohl A, Tositti L, Tubertini O, Colombo T: Stratospheric ozone intrusion episodes recorded at Mt.Cimone during the VOTALP project: case studies, Atmos. Environ., 34 (9), 2000.

<http://www.sciencedirect.com/science/article/pii/S1352231099002800>

Seibert P, Feldmann H, Neininger B, Baumle M, Trickl T: South foehn and ozone in the Eastern Alps: case study and climatological aspects, Atmos. Environ., 34 (9), 2000.

<http://www.sciencedirect.com/science/article/pii/S1352231099004392>

Balkanski Y, Schulz M, Bonasoni P, van Dingenen R, Hanke M, Gobbi G, Kormann R, Calzolari F: Measurements of Aerosol, Chemically Active Species Properties and Evidence for Their Interactions During the Mt Cimone (ITALY) Campaign: June 1st 2000 - July 6th 2000, F. Eos Trans, 81 (48), 2000.

Stohl A, Spichtinger-Rakowsky N, Bonasoni P, Feldmann H, Memmesheimer M, Scheel HE, Trickl T, Hübener SH: The influence of stratospheric intrusions on alpine ozone concentrations, Atmos. Environ., 34 (9), 2000.

<http://www.sciencedirect.com/science/article/pii/S1352231099003209>

Colombo T, Santaguida R, Capasso A, Calzolari F, Evangelist F, Bonasoni P: Biospheric influence on carbon dioxide measurements in Italy. Atmos. Environ., 34 (29-30), 2000.

<http://www.sciencedirect.com/science/article/pii/S1352231000003666>

Mangani F, Maione M, Lattanzi L, Arduini J: Atmospheric measurements of the halogenated hydrocarbons involved in global change phenomena, Atmos. Environ., 34 (29-30), 2000.

<http://www.sciencedirect.com/science/article/pii/S1352231000002247>

Gerasopoulos E, Zanis P, Stohl A, Zerefos CS, Papastefanou C, Ringerd W, Tobler L, Hubener S, Gaeggler HW, Kanter HJ, Tositti L, Sandrini S: A climatology of ^{7}Be at four high-altitude stations at the Alps and the Northern Apennines, Atmos. Environ., 35, 2001.

<http://www.sciencedirect.com/science/article/pii/S1352231001004009>

Tomasi C, S. Marani, V. Vitale, Lupi A: Convective transport of particulate matter in an Apennine valley - 2. Time variations in the columnar aerosol mass content and vertical profiles of aerosol mass concentration, Atmos. Res., 63, 2002.

<http://www.sciencedirect.com/science/article/pii/S0169809502000376>

Marani S, Tomasi C, Vitale V: Convective transport of particulate matter in an Apennine valey 1. Sun-photometric measurements of aerosol optical depth time-variations at various altitudes, Atmos. Res., 61. 2002.

<http://www.sciencedirect.com/science/article/pii/S016980950100103X>

Hanke M, Uecker J, Reiner T, Arnold F: Atmospheric peroxy radicals: ROXMAS, a new massspectrometric methodology for speciated measurements of HO_2 and SRO_2 and first results, Int. J. Mass. Spectrum., 213, 2002.

Kormann R, Fischer H, Gurk C, Helleis F, Klupfel T, Kowalski K, Konigstedt R, Parchatka U, Wagner V: Application of a multi-laser tunable diode laser absorption spectrometer for atmospheric trace gas measurements at sub-ppbv levels, Spectrochim. Acta.,Part A, 58 (2489), 2002.

<http://www.sciencedirect.com/science/article/pii/S1386142502000665>

Stohl A, Bonasoni P, Cristofanelli P, Collins W, Feichter J, Frank A, Forster C, Gerasopoulos E, Gäggeler H, James P et al: Stratosphere-troposphere exchange - a review, and what we have learned from STACCATO, J. Geophys. Res., 108

(D12), 2003.

<http://onlinelibrary.wiley.com/doi/10.1029/2002JD002490/pdf>

Cristofanelli P, Bonasoni P, Collins W, Feichter J, Forster C, James P, Kentarchos A, Kubik PW, Land C, Meloen J: Stratosphere-to-troposphere transport: A model and method evaluation, *J. Geophys. Res.*, 108 (D12), 2003.

<http://onlinelibrary.wiley.com/doi/10.1029/2002JD002600/full>

Fischer H, Kormann R, Klupfel T, Gurk C, Konigstedt R, Parchatka U, Muhle J, Rhee TS, Brenninkmeijer CAM, Bonasoni P, Stohl A: Ozone production and trace gas correlations during the June 2000 MINATROC intensive measurement campaign at Mt. Cimone., *Atmos. Chem. Phys.*, 3, 2003.

<http://www.atmos-chem-phys.net/3/725/2003/acp-3-725-2003.pdf>

Balkanski Y, Bauer SE, van Dingenen R, Bonasoni P, Schulz M, Fischer H, Gobbi GP, Hanke M, Hauglustaine D, Putaud JP, Stohl A, Raes F: The Mt Cimone, Italy, free tropospheric campaign: principal characteristics of the gaseous and aerosol composition from European pollution, Mediterranean influences and during African dust events, *Atmos. Chem. Phys. Disc.*, 3, 2003.

<http://www.atmos-chem-phys-discuss.net/3/1753/2003/acpd-3-1753-2003.pdf>

Putaud J-P, Dingenen VR, Dell'Acqua A, Raes F, Matta E, Decesari S, Facchini MC, Fuzzi S: Size-segregated aerosol mass closure and chemical composition in Monte Cimone (I) during MINATROC, *Atmos. Chem. Phys.*, 3, 2003.

<http://www.atmos-chem-phys.net/4/889/2004/acp-4-889-2004.pdf>

Zanis P, Trickl T, Stohl A, Wernli H, Cooper O, Zerefos C, Gaeggeler H, Schnabel C, Tobler L, Kubik PW et al.: Forecast, observation and modelling of a deep stratospheric intrusion event over Europe, *Atmos. Chem. Phys.*, 3, 2003.

<http://www.atmos-chem-phys.net/3/763/2003/acp-3-763-2003.pdf>

Hanke M, Umann B, Uecker J, Arnold F, Bunz H: Atmospheric measurements of gas-phase HNO₃ and SO₂ using chemical ionization mass spectrometry during the MINATROC field campaign 2000 on Monte Cimone, *Atmos. Chem. Phys.*, 3, 417–436, 2003.

<http://www.atmos-chem-phys.net/3/417/2003/acp-3-417-2003.pdf>

A. Petritoli, Bonasoni P, Giovanelli G, Ravagnani F, Kostadinov I, Bortoli D, Weiss A, Schaub D, Richter A, Fortezza F: First Comparison Between ground-based and Satellite-borne Measurements of Tropospheric Nitrogen Dioxide in the Po Basin, *J. Geophys. Res.*, 109 (D15), 2004.

<http://onlinelibrary.wiley.com/doi/10.1029/2004JD004547/full>

Bonasoni P, Cristofanelli P, Calzolari F, U. Bonafe, Evangelisti F, Stohl A, Zauli Sajani S, van Dingenen R, Colombo T, Balkanski Y: Aerosol-ozone correlations during dust transport episodes, *Atmos. Chem. Phys.*, 4, 2004.

<http://www.atmos-chem-phys.net/4/1201/2004/acp-4-1201-2004.pdf>

Van Dingenen R, Putaud JP, Martins-Dos Santos S, Raes F: Physical aerosol properties and their relation to air mass origin at Monte Cimone (Italy) during the first MINATROC campaign, *Atmos. Chem. Phys.*, 5, 2203–2226, 2005.

<http://www.atmos-chem-phys.net/5/2203/2005/acp-5-2203-2005.pdf>

Campana M, Li Y, Staehelin J, Prévôt ASH, Bonasoni P, Loetscher HP, Peter T: The influence of south Foehn on the ozone mixing ratios at the alpine site Arosa, *Atmos. Environm.*, 39, 2005.

<http://www.sciencedirect.com/science/article/pii/S1352231005000476>

Beine HJ, Amoroso A, Esposito G, Sparapani R, Ianniello A, Georgiadis T, Nardino M, Bonasoni P, Cristofanelli P, Domine' F: Deposition of atmospheric nitrous acid on alkaline snow surfaces, *Geophys. Res. Lett.*, 32 (L10808), 2005.

<http://onlinelibrary.wiley.com/doi/10.1029/2005GL022589/full>

Greally BR, Manning AJ, Reimann S, McCulloch A, Huang J, Dunse BL, Simmonds PG, Prinn RG, Fraser PJ, Cunnold DM et al.: Observation of 1,1-difluoroethane (HFC-152a) at AGAGE and SOGE monitoring stations 1994-2004 and derived Global and regional emission estimates, *J. Geophys. Res.*, 112, 2006.

<http://onlinelibrary.wiley.com/doi/10.1029/2006JD007527/pdf>

Marenco F, Bonasoni P, Calzolari F, Ceriani M, Chiari M, Cristofanelli P, D'Alessandro A, Fermo P, Lucarelli F, Mazzei F et al.: Characterization of atmospheric aerosols at Monte Cimone, Italy, during summer 2004: source apportionment and transport mechanisms, *J. Geophys. Res.*, 111 (D24202), 2006.

<http://onlinelibrary.wiley.com/doi/10.1029/2006JD007145/full>

Cristofanelli P, Bonasoni P, Tositti L, Bonafe' U, Calzolari F, Evangelisti F, Sandrini S, Stohl A: A 6-year analysis of stratospheric intrusions and their influence on ozone at Mt. Cimone (2165 m above sea level), *J. Geophys. Res.*, 111

(D03306), 2006.

<http://onlinelibrary.wiley.com/doi/10.1029/2005JD006553/full>

Colombo T, Pelino V, Vergari S, Cristofanelli P, Bonasoni P: Study of temperature and precipitation variations in Italy based on surface instrumental observations, *Global Planet. Change*, 57 (3-4), 2007.

<http://www.sciencedirect.com/science/article/pii/S0921818106003250>

Cristofanelli P, Bonasoni P, Carboni G, Calzolari F, Casarola L, Sajani ZS, Santaguida R: Anomalous high ozone concentrations recorded at a high mountain station in Italy in Summer 2003., *Atmos. Env.* 41, 2007.

<http://www.sciencedirect.com/science/article/pii/S1352231006010326>

Lee HN, Tositti L, Zheng XD, Bonasoni P: Analyses and comparisons of ^{7}Be , ^{210}Pb and activity ratio $^{7}\text{Be}/^{210}\text{Pb}$ with ozone observations at two GAW stations from high mountains, *J. Geophys. Res.*, 112 (D05303), 2007.

<http://onlinelibrary.wiley.com/doi/10.1029/2006JD007421/pdf>

Cristofanelli P, Calzolari F, Bonafè U, R.Duchi, Marinoni A, Roccato F, Tositti L, Bonasoni P: Stratospheric Intrusion Index (SI2) from baseline measurement data, *Theor. App. Clim.*, 2008.

<http://link.springer.com/article/10.1007/s00704-008-0073-x>

Maione M, Giostra U, Arduini J, Belfiore L, Furlani F, Geniali A, Mangani G, Vollmer MK, Reimann S: Localization of source regions of selected hydrofluorocarbons combining data collected at two European mountain Stations, *Sci. Total Environ.*, 391, 232–240, 2008.

<http://www.sciencedirect.com/science/article/pii/S0048969707010832>

Marinoni A, Cristofanelli P, Calzolari F, Roccato F, Bonafe' U, Bonasoni. P.: Continuous measurements of aerosol physical parameters at the Mt. Cimone GAW Station (Italy - 2165 m a.s.l.), *Sci. Total Environ.*, 391, 2008.

<http://www.sciencedirect.com/science/article/pii/S0048969707010844#>

Cristofanelli P, Bonasoni P: Background ozone in the southern Europe and Mediterranean area: influence of the transport processes, *Environ. Poll.*, 2008.

<http://www.sciencedirect.com/science/article/pii/S026974910800451X>

Cristofanelli P, Marinoni A, Arduini J, Bonafè U, Calzolari F, Colombo T, Decesari S, Duchi R, Facchini MC, Fierli F et al.: Significant variations of trace gas composition and aerosol properties at Mt. Cimone during air mass transport from North Africa – contributions from wildfire emissions and mineral dust, *Atmos. Chem. Phys.*, 9, 2009.

<http://www.atmos-chem-phys.net/9/4603/2009/acp-9-4603-2009.pdf>

Stohl A, Seibert P, Arduini J, Eckhardt S, Fraser P, Greatly BR, Maione M, O'Doherty S, Prinn RG, Reimann S et al.: A new analytical inversion method for determining regional and global emissions of greenhouse gases: sensitivity studies and application to halocarbons, *Atmos. Chem. Phys.*, 9, 1597–1620, 2009.

<http://www.atmos-chem-phys.net/9/1597/2009/acp-9-1597-2009.pdf>

Muller T, et al.: Angular Illumination and Truncation of Three Different Integrating Nephelometers: Implications for Empirical, Size-Based Corrections. *Aerosol Sci. Tech.*, 43 (6), 2009.

<http://www.tandfonline.com/doi/full/10.1080/02786820902798484#.UlfM41O9KuY>

Carbone C, Decesari S, Mircea M, Giulianelli L, Finessi E, Rinaldi M, Fuzzi S, Marinoni A, Duchi R, Perrino C et al.: Size-resolved aerosol chemical composition over the Italian Peninsula during typical summer and winter conditions, *Atmos. Environ.*, 44 (39), 5269-5278, 2010.

<http://www.sciencedirect.com/science/article/pii/S1352231010006618>

Sajani ZS, Miglio R, Bonasoni P, Cristofanelli P, Marinoni A, Sartini C, Goldoni CA, Girolamo DG, Lauriola P: Saharan dust and daily mortality in Emilia-Romagna (Italy), *Occup Environ Med*, 2010.

<http://oem.bmj.com/content/68/6/446.full.pdf+html>

Xiao X, Prinn RG, Fraser PJ, Simmonds PG, Weiss RF, O'Doherty S, Miller BR, Salameh PK, Harth CM, Krummel PB et al.: Optimal Estimation of the Surface Fluxes of Methyl Chloride using a 3-D Global Chemical Transport Model, *Atmos. Chem. Phys.*, 10, 5515-5533, 2010-

<http://www.atmos-chem-phys.net/10/5515/2010/acp-10-5515-2010.pdf>

Petkov B, Tomasi C, Vitale V, di Sarra A, Bonasoni P, Lanconelli C, Benedetti E, Sferlazzo D, Diemoz H, Agnesod G et al.: Ground-based observations of solar radiation at three Italian sites, during the eclipse of 29 March, 2006: Signs of the environment impact on incoming global irradiance, *Atmos. Res.*, 96 (1), 2010.

<http://www.sciencedirect.com/science/article/pii/S0169809509003457>

Asmi A, et al. : Number size distributions and seasonality of submicron particles in Europe 2008–2009, *Atmos. Chem. Phys.*, 11, 2011.

<http://www.atmos-chem-phys.net/11/5505/2011/acp-11-5505-2011.pdf>

Yver CE, et al. : A new estimation of the recent tropospheric molecular hydrogen budget using atmospheric observations and variational inversion, *Atmos. Chem. Phys.*, 11, 2011.

<http://www.atmos-chem-phys.net/11/3375/2011/acp-11-3375-2011.pdf>

Keller CA, Hill M, Vollmer M. K., Henne S, Brunner D, Reimann S, O'Doherty S, Arduini J, Maione M, Ferenczi Z et al.: European Emissions of Halogenated Greenhouse Gases Inferred from Atmospheric Measurements, *Environ. Sci and Technol.*, 46 (1), 217-225, 2011.

<http://pubs.acs.org/doi/10.1021/es202453j>

Yver CE et al.: A new estimation of the recent tropospheric molecular hydrogen budget using atmospheric observations and variational inversion, *Atmos. Chem. and Phys.*, 11, 2011.

<http://www.atmos-chem-phys.net/11/3375/2011/acp-11-3375-2011.html>

Giostra U, Furlani F, Arduini J, Cava D, Manning AJ, O'Doherty SJ, Reimann S, Maione M: The determination of a regional atmospheric background mixing ratio for anthropogenic greenhouse gases: a comparison of two independent methods, *Atmos. Environ.*, 45, 2011.

<http://www.sciencedirect.com/science/article/pii/S135223101100700X>

Andrews E, et al. : Climatology of Aerosol Radiative Properties in the Free Troposphere, *Atmos. Res.*, 102, 2011.

<http://www.sciencedirect.com/science/article/pii/S0169809511002857>

Muller T, et al.: Characterization and intercomparison of aerosol absorption photometers: result of two intercomparison workshops, *Atmos. Meas. Tech.*, 4, 2011.

<http://www.atmos-meas-tech.net/4/245/2011/amt-4-245-2011.pdf>

Abeli T, Rossi G, Gentili R, Mondoni A, Cristofanelli P: Response of alpine plant flower production to temperature and snow cover fluctuation at the species range boundary, *Plant Ecol.*, 213 (1 (2012)), 1-13, 2012.

<http://link.springer.com/article/10.1007%2Fs11258-011-0001-5>

Zauli-Sajani S, P.Bonasoni, Cristofanelli P, Marinoni A, Lauriola P: Only coarse particles from the Sahara?, *Epidemiology*, 4(23), 2012.

<http://journals.lww.com/epidem/pages/results.aspx?txtKeywords=%22Sajani%22>

Abeli T, Rossi G, Gentili R, Gandini M, Mondoni A, Cristofanelli P: Effect of the extreme summer heat waves on isolated populations of two orophytic plants in the north Apennines (Italy), *Nordic J. Bot.*, 1 (30), 2012.

<http://onlinelibrary.wiley.com/doi/10.1111/j.1756-1051.2011.01303.x/abstract>

Saikawa E et al.: Global and regional emissions estimates for HCFC-22, *Atmos. Chem. Phys.*, 12, 2012

<http://www.atmos-chem-phys.net/12/10033/2012/acp-12-10033-2012.html>

Brunner D, Henne S, Keller CA Reimann S, Vollmer MK, O'Doherty S, and Maione M: An extended Kalman-filter for regional scale inverse emission estimation, *Atmos. Chem. Phys.*, 12, 3455-3478, doi:10.5194/acp-12-3455-2012, 2012

<http://www.atmos-chem-phys.net/12/3455/2012/acp-12-3455-2012.html>

Tositti L, Riccio A, Sandrini S, Brattich E, Baldacci D, Parmeggiani S, Cristofanelli P, Bonasoni P: Short-term climatology of PM10 at a high altitude background station in southern Europe, *Atmos. Environ.*, 65, 2013.

<http://www.sciencedirect.com/science/article/pii/S1352231012010333>

Weaver C, Kiemle C, Kawa SR, Aalto T, Necki J, Steinbacher M, Arduini J, Apadula F, Berkhouit H, Hatakka J et al.: Retrieval of methane source strengths in Europe using a simple modeling approach to assess the potential of space-borne lidar observations, *Atmos. Chem. Phys. Discuss.*, 13, 2013.

<http://www.atmos-chem-phys-discuss.net/13/19559/2013/acpd-13-19559-2013.html>

Cristofanelli P, Fierli F, Marinoni A, Calzolari F, Duchi R, Burkhardt J, Stohl A, Maione M, Arduini J, Bonasoni P: Influence of biomass burning and anthropogenic emissions on ozone, carbon monoxide and black carbon at the Mt. Cimone GAW-WMO global station (Italy, 2165 m a.s.l.), *Atmos. Chem. Phys.*, 13, 2013.

<http://www.atmos-chem-phys.net/13/15/2013/acp-13-15-2013.pdf>

Cristofanelli P, di Carlo P, Altorio AD, Salisburgo DC, Tuccella P, Biancofiore F, Stocchi P, Verza GP, Landi TC, Marinoni A et al.: Analysis of Summer Ozone Observations at a High Mountain Site in Central Italy (Campi Imperatore, 2388 m a.s.l.), Pure and Appl. Geophys., 2013.

<http://link.springer.com/article/10.1007/s00024-012-0630-1>

Maione M, Giostra U, Arduini J, Furlani F, Graziosi F, Lo Vullo E, and Bonasoni P: Ten years of continuous observations of stratospheric ozone depleting gases at Monte Cimone (Italy) - Comments on the effectiveness of the Montreal Protocol from a regional perspective, Sci. Tot. Environ., 445–446, 2013.

<http://www.sciencedirect.com/science/article/pii/S0048969712016130>

Hall BD et al.: Results from the International Halocarbons in Air Comparison Experiment (IHALACE), Atmos. Meas. Tech., 6, 2013.

<http://www.atmos-meas-tech.net/7/469/2014/amt-7-469-2014.html>

Tositti L, Brattich E, Cinelli G, Baldacci D.: 12 years of ^7Be and ^{210}Pb data at the WMO-GAW station of Mt. Cimone (2165 m a.s.l., 44°12'N 10°42'E) and their correlation with meteorological parameters. Atmos. Environ., 2014.

Carbone C, De Cesari S, Paglione M, Julianelli L, Rinaldi M, Marinoni A, Cristofanelli P, Didiodato A.: 3-year chemical composition of free tropospheric PM1 at the Mt. Cimone GAW global station – South Europe – 2165 m a.s.l., Atmos. Environ., 87, 2014.

Sandrin S, et al.: Spatial and seasonal variability of carbonaceous aerosol across Italy, Atmos. Environ., 99, 2014

Cristofanelli P, Scheel H-E, Steinbacher M, Saliba M., et al.: Long-term surface ozone variability at Mt. Cimone WMO/GAW global station (2165 m a.s.l., Italy. Atmos. Environ., 101:23-33, 2015.

Brattich E, Riccio A, Tositti L, Cristofanelli P, Bonasoni P.: An outstanding Saharan dust event at Mt. Cimone (2165 m a.s.l., Italy) in March 2004, Atmos. Environ., 113, 2015.

Rinaldi M, Gilardoni S, Paglione M, Sandrin S, Fuzzi S, Massoli P, Bonasoni P, Cristofanelli P, Marinoni A, Poluzzi V et al.: Organic aerosol evolution and transport observed at Mt. Cimone (2165 m a.s.l.), Italy, during the PEGASOS campaign. Atmos. Chem. Phys., 15, 2015.

Brattich E, Hernández-Ceballos MA, Cinelli G, Tositti L.: Analysis of ^{210}Pb peak values at Mt. Cimone (1998–2011), Atmos. Environ., 112, 2015.

Research Projects



GAW (Global Atmosphere Watch) The Global Atmosphere Watch (GAW) programme of WMO is a partnership involving 80 countries, which provides reliable scientific data and information on the chemical composition of the atmosphere, its natural and anthropogenic change, and helps to improve the understanding of interactions between the atmosphere, the oceans and the biosphere.



NextData The Project of Interest NextData will favour the implementation of measurement networks in remote mountain and marine areas and will develop efficient web portals to access meteoclimatic and atmospheric composition data, past climate information from ice and sediment cores, biodiversity and ecosystem data, measurements of the hydrological cycle, marine reanalyses and climate projections at global and regional scale.



SHARE (Station at High Altitude for Research on the Environment) Share is an integrated project funded by EV-K2-CNR Committe comprising an international climate and atmospheric monitoring network, researches in environmental and geophysical sciences and new technology development for monitoring activity in high mountain regions. Working in synergy with projects run by UNEP and WMO, data from the SHARE initiative benefit the international scientific community as well as decision makers.



ACTRIS (Aerosols, Clouds, and Trace gases Research InfraStructure Network) ACTRIS is a European Project aiming at integrating European ground-based stations equipped with advanced atmospheric probing instrumentation for aerosols, clouds, and short-lived gas-phase species. ACTRIS will have the essential role to support building of new knowledge as well as policy issues on climate change, air quality, and long-range transport of pollutants. ACTRIS is building the next generation of the ground-based component of the EU observing system by integrating three existing research infrastructures EUSAAR, EARLINET, CLOUDNET, and a new trace gas network component into a single coordinated framework. ACTRIS is funded within the EC 7th Framework Programme under "Research Infrastructures for Atmospheric Research" and started on 1 April 2011 for a period of 4 years.



MACC-2 (Monitoring Atmospheric Composition and Climate - Interim Implementation) is the current pre-operational atmospheric service of the European GMES programme. MACC provides data records on atmospheric composition for recent years, data for monitoring present conditions and forecasts of the distribution of key constituents for a few days ahead. MACC-II combines state-of-the-art atmospheric modelling with Earth observation data to provide information services covering European Air Quality, Global Atmospheric Composition, Climate, and UV and Solar Energy.



EUSAAR (European Supersites for Atmospheric Aerosol Research) The objective of EUSAAR UE-funded project is the integration of measurements of atmospheric aerosol properties performed in a distributed network of 20 high quality European ground-based stations. This integration contributes to a sustainable reliable operational service in support of policy issues on air quality, long-range transport of pollutants and climate change.



EUROHYDROS The aim of EUROHYDROS has been the initialisation of a European Network for observations of molecular Hydrogen based on 12 continuous measurements sites which allow a wide range of observation, from clean air stations for measurements of atmospheric background to moderately polluted and urban. This in order to improve the understanding of hydrogen in the global background atmosphere and of the impact of European emissions on the present day atmosphere.



CIRCE (Climate Change and Impact Reserach: the Mediterranean Environment) The general project objectives are to predict and to quantify physical impacts of climate changes in the Mediterranean area; to evaluate the consequences of climate changes for the society and the economy of the populations located in the Mediterranean area; to develop an integrated approach to understand combined effects of climate change; to identify adaptation and mitigation strategies in collaboration with regional stakeholders.



AGAGE-Advanced Global Atmospheric Gases Experiment AGAGE and its predecessors (the Atmospheric Life Experiment, ALE, and the Global Atmospheric Gases Experiment, GAGE) have been measuring the composition of the global atmosphere continuously since 1978. The AGAGE is distinguished by its capability to measure over the globe at high frequency almost all of the important gases species in the Montreal Protocol (e.g. CFCs and HCFCs) to protect the ozone layer and almost all of the significant non-CO₂ gases in the Kyoto Protocol (e.g. HFCs, methane, and nitrous oxide) to mitigate climate change.



CEOP HE (Coordinated Energy and Water Cycle Observation Project - High Elevation) CEOP HE is a component of 'regional focus' within the Coordinated Energy and Water Cycle Observation Project (CEOP) of the Global Energy and Water Cycle Experiment (GEWEX), under the WCRP of WMO. CEOP HE aims to further knowledge on physical and dynamic processes in high elevation areas, contributing to global climate and water cycle studies by providing rare but crucial information from high elevations. This initiative was launched and is coordinated by the Ev-K2-CNR Committee.



ACCENT
ATMOSPHERIC COMPOSITION CHANGE
THE EUROPEAN NETWORK OF EXCELLENCE

ACCENT (Atmospheric Composition Change - The European Network of Excellence) The overall goals of the UE-network ACCENT are to promote a common European strategy for research on atmospheric composition sustainability, to develop and maintain durable means of communication and collaboration within the European scientific community, to facilitate this research and to optimise the interactions with policy-makers and the general public.

AEROCLOUDS (Climatic Effects of Aerosol and Clouds) AEROCLOUDS is a project funded by the Italian Ministry for University and Research to improve our knowledge of the role of aerosol and clouds in the climate system. Four research lines have been investigated: 1) Radiative properties of aerosols ("direct" climatic effects); 2) Aerosol-Cloud interactions ("indirect" climatic effects); 3) Climatic effects of clouds and precipitation; 4) Regional and global modelling of the aerosol effects on climate.



SOGE (System for Observations of Halogenated Greenhouse Gases in Europe) SOGE is an integrated system for observation of halogenated greenhouse gases in Europe. The project was funded by UE and builds on a combination of observations and modelling. The observations are partly surface in situ data collected continuously at four background stations as a part of national or international measurement programs. For some species(PFC, SF₆), for which high-frequency measurements are not yet fully developed, such capacity will be developed as a part of SOGE.



POLPO (Pollution Hot Spot Monitoring from GOME Applied to the Po-basin) POLPO investigated the feasibility of applying satellite data for monitoring large pollution plumes. The prototype service demonstrated the application of GOME data for case studies. Users as, e.g., environmental agencies, who have to rely on ground-based measurements, found the added value satellited data provide together with its limitations in the feasability study.

QUILT

QUILT (Quantification and Interpretation of Long-Term UV-Vis Observations of the Stratosphere)

The general aim was to use the existing ground-based, satellite and balloon borne UV-visible data as well as 3D atmospheric modelling tools for quantifying ozone loss in the past, to monitor its development in the present and to investigate its relation to active halogen and nitrogen species.



TOR-2 (Tropospheric Ozone Research - 2) The overall aim of TOR-2 was to quantify crucial processes in the atmosphere in order to improve the scientific background for the development of effect-based control strategies for photochemical oxidants over Europe.



STACCATO (Influence of Stratosphere-Troposphere Exchange in a Changing Climate on Atmospheric Transport and Oxidation Capacity) EU-project STACCATO is a comprehensive study of stratosphere-troposphere exchange (STE) processes and their effect on atmospheric chemistry. STE is a key factor controlling the budget of ozone, water vapour and other substances in both the troposphere and lower stratosphere.



MINATROC (Mineral Dust and Tropospheric Chemistry) Problems to be solved this EU-project focuses on the transformation of atmospheric pollutants from Europe in the presence of mineral dust over South Europe and Africa. Intensive field campaigns, experimental laboratory investigations and modeling studies were conducted to evaluate the influence of mineral dust on tropospheric oxidizing properties.



VOTALP-2 (Vertical Ozone Transport in the Alps - 2) The EU research project VOTALP II investigated the enhanced vertical exchange above the Alps as well as other processes which might be relevant for increased ozone concentrations. The role of stratospheric intrusions for mountain peaks and of horizontal advection of polluted air for the foothill area causing a high ozone abundance has been quantified for selected locations.



VOTALP (Vertical Ozone Transport in the Alps) The EU research project VOTALP investigated transport and formation of ozone in the Alps, focusing on processes which can cause increased ozone concentrations, namely stratospheric intrusions, horizontal advection of polluted air, and in-situ production of ozone due to emissions in Alpine valleys.

The ICO-OV staff

Paolo Bonasoni – *Station leader*
p.bonasoni@isac.cnr.it

Paolo Cristofanelli – *Trace gases and meteorology*
p.cristofanelli@isac.cnr.it

Angela Marinoni – *Atmospheric aerosol*
a.marinoni@isac.cnr.it

Francescopiero Calzolari – *Data acquisition and technical infrastructures*
f.calzolari@isac.cnr.it

Michela Maione (Urbino University) - *VOC and Halogenated gases*
michela.maione@uniurb.it

Jgor Arduini (Urbino University) - *VOC and Halogenated gases*
jgor.arduini@uniurb.it

Fabrizio Roccato – *System manager*
f.roccato@isac.cnr.it

Rocco Duchi – *Data validation and analysis*
r.duchi@isac.cnr.it

Davide Putero – *Data analysis*
d.putero@isac.cnr.it

Tony Christian Landi – *Data analysis*
t.landi@isac.cnr.it

Ubaldo Bonafè - *Technical infrastructures*
u.bonafe@isac.cnr.it

CONTACTS:

Italian Climate Observatory “O. Vittori” at Mt. Cimone
c/o Institute for Atmospheric Sciences and Climate – National Research Council of Italy
Via Gobetti 101
40129 Bologna (Italy)
www.isac.cnr.it/cimone





<http://www.isac.cnr.it/cimone/>